

PSORIASIS SPECIALTY CARE PROGRAM

Phone: **844-378-3784** • Fax: **888-972-7058**



PATIENT IN			2 PRESCRIBER INFORMATION: Name:				
vame							
Citv:	State: Ziµ	City:	City: State: Zip:				
Phone:	Alt. Phone:	Priorie:	Priorie: Fax: Fax:				
mail:		NET	NET DEA				
OOB:	Gender: O M O F Caregiver	Tax I D ·	Supervising Physician NPI Tax I.D.: Tax I.D.: NPI				
leight:	Weight: Allergies:		Office Contact: P		Phone:		
_	T OF MEDICAL NECESSIT	(Please Attach All Medical Documentation) tient also taking Methotrexate? □ Yes □ No		Prior Failed Treatments: ☐ Topicals			
ICD-10: TB Test: □ Positive LFT: ALT: Assessment: □ Moo % BSA affecto □ Scalp □ Face □	Other: Negative Date: AST: Date: derate \(\Boxed{\text{Mod to Severe}} \) Severe	Serious or active infection pre Hep B ruled out or treatment so Does patient have latex allergy If Prior Authorization is denied, re formulary alternatives will be pro- based upon the patient's insurar	☐ Methotrexate ☐ Oral Meds ☐ Biologics ☐ UVA ☐ UVB ☐ Others				
3 INJECTION	TRAINING: O To be Administered by a	ealthcare Provider O Pharmacist to P	rovide Training O Patient Trair	ed in MD Office O Manufactu	ırer Nurse	Support	
3 PRODUCT	DELIVERY: O Patient's Hom	Physician's Office	O Pharmacy to Coo	ordinate			
PRESCRIPTION	INFORMATION: Please Inclu- INFORMATION: (Please be s	re to choose both induct	on and maintenance		-		
Medication	Dosage & Strength		Direction		QTY	Refills	
☐ CIMZIA®	☐ 200mg/ml Prefilled Syringe☐ 200mg Lyophilized Powder Vial		y other week ight <90kg) Inject 400mg SC i (Weight <90kg) Inject 200mg i	-			
☐ COSENTYX®	☐ 150mg/ml Sensoready® Pen☐ 150mg/ml Prefilled Syringe	☐ Induction Dose: Inject 150mg SC at weeks 0, 1, 2, 3, and 4☐ Induction Dose: Inject 300mg SC at weeks 0, 1, 2, 3, and 4☐ Ind		5 10	0		
	□ 150mg/ml Lyophilized Powder Vial	☐ Maintenance Dose: Inject 150mg SC every four weeks☐ Maintenance Dose: Inject 300mg SC every four weeks			1 2		
☐ ENBREL®	□ 50mg/ml Sureclick Autoinjector		duction Dose: Inject 50mg SC twice a week 4 days apart) for 3 months, then start maintenance dosing			2	
	☐ 50mg/ml Enbrel Mini™ Prefilled Cartridge	For		4			
	Enbrel Mini™ only: AutoTouch™ Autoinj		Pediatric Patients: To achieve pediatric doses other than				
	□ 50mg/ml Prefilled Syringe□ 25mg/0.5ml Prefilled Syringe	9	50mg or 25mg, use reconstituted Enbrel lyophilized powder > 63 kg or more: Inject 50mg weekly < 63 kg: Inject 0.8mg/kg weekly Other:		4		
	□ 25mg Lyophilized Powder Multiple Dose □ Other:	ui					
	- Juigi.	Psoriasis					
	☐ Psoriasis 80mg/0.8ml and 40mg/0.4ml	☐ Induction Dose: Inje	ct 80mg SC on day 1, then 40 SC every other week	mg SC	3	0	
	Starter Package ☐ Psoriasis 40mg/0.4ml Starter Package	•	•		4	0	
	☐ 40mg/0.4ml Pen	Other:	■ Maintenance: Inject 40mg SC every other week ■ Other:		2		
	□ 80mg/0.8ml Pen □ 40mg/0.4ml Prefilled Syringe □ 80mg/0.8ml Prefilled Syringe	Psoriatic Arthritis ☐ Inject 40 mg SC eve	Psoriatic Arthritis ☐ Inject 40 mg SC every other week				
□ HUMIRA®	Hidradenitis Suppurative	Induction Dose:				0	
	 ☐ Hidradenitis Suppurativa 80mg/0.8ml Starter Package ☐ Hidradenitis Suppurativa 40mg/0.4ml 	☐ Inject one 80mg pen S				0	
	Starter Package 40mg/0.4ml Pen 40mg/0.8ml Pen 80mg/0.8ml Pen 40mg/0.4ml Prefilled Syringe 80mg/0.8ml Prefilled Syringe	Maintenance: ☐ Inject 40mg SC on day 29 and every week thereafter ☐ Inject 80mg SC on day 29 and every other week thereafter ☐ Patient has signed HUMIRA Complete form			4		
<u> </u>	All stren	hs and dosages listed are Humira® (nuale rree				
	R SIGNATURE: I authorize pharmacy	act as my designee for initiating and coordii	nating insurance prior authorizations	s, nursing services and patient ass	 istance pro	grams.	
Signature:		e: Signature	Dispense As \	Date			



PSORIASIS SPECIALTY CARE PROGRAM

Phone: **844-378-3784** • Fax: **888-972-7058**



v10.8_081721

©2020 KloudScript, Inc. - All rights reserved.

PATIENT IN			PRESCRIBER INFORI			
Name:			Name: Address:			
Address	Stato: 7ii		City: State: Zip:			
	State: Zip: Alt. Phone:		Phone:	Fax:		
Friorie Email:	Alt. Filone		NPI:	_ DEA:		
LITIAII	Gender: O M O F Caregiver:		Supervising Physician:	NPI:		
оов	_ delider. 9 ivi 91 Galegiver		- Tax I.D.: Phone: Phone:			
				Priorie:		
3 STATEMENT OF MEDICAL NECESSITY: (Please Attach A Date of Diagnosis: Patient also taki			Medical Documentation) ∫ Methotrexate? □ Yes □ No	Prior Failed Treatments: ☐ Topicals		
ICD-10:	Other: Serious or active		nfection present? ☐ Yes ☐ No	☐ Methotrexate		
TB Test: ☐ Positive	Pate: Hep B ruled out		or treatment started? ☐ Yes ☐ No ☐ Oral Meds			
	_ AST: Date: Thep b rated of the derate ☐ Mod to Severe ☐ Severe Does patient h		latex allergy? ☐ Yes ☐ No	_		
% BSA affected		If Prior Authorization is denied, recommended				
☐ Scalp ☐ Face ☐ Chest ☐ Arms ☐ Hands ☐ Nails		formulary alternatives will be provided to the prescriber UVA UVB				
☐ Back ☐ Groin ☐	□ Back □ Groin □ Buttocks □ Legs □ Other: based upon the		ne patient's insurance coverage.			
-	TRAINING: O To be Administered by a				rer Nurse	Support
5 PRODUCT	DELIVERY: O Patient's Hom	ne O Physiciar	n's Office O Pharmacy to Co	ordinate		
	INFORMATION: Please Inclu- INFORMATION: (Please be s		' '	e dose where applica	able)	
Medication	Dosage & Strength		Direction		QTY	Refills
☐ ORENCIA®	☐ 125mg/ml ClickJect [™] Autoinjector☐ 125mg/ml Prefilled Syringe	☐ Inject 12	5mg SC once a week		4	
☐ OTEZLA®	- otation rack (milation)		er Pack: Take one tablet in the morning on day 1, then take one tablet in orning and one tablet in the evening as directed on the starter pack		1	0
	□ 30mg Tablets □ Maint		enance: Take one 30mg tablet by mouth twice daily			
	☐ 17.5mg ☐ 10mg ☐ 12.5mg ☐ 15mg ☐ 17.5mg ☐ 20mg ☐ 22.5mg ☐ 25mg ☐ *An initial		mg SC once weekly			
☐ RASUVO®			al test dose of 2.5 to 5 mg is recommended in patients with risk or hematologic toxicity or renal impairment*			
SIMPONI® (for PSA)	☐ 50mg/0.5ml Prefilled syringe ☐ 50mg/0.5ml Prefilled SmartJect® autoinje	□ Inject 50	mg SC once a month		1	
			on Dose: Inject 150mg (two 75mg injections) SC at weeks 0 and 4	4	0
□ SKYRIZI [™]	☐ 150mg/ml in each single-dose prefilled s	. u iviainten	enance Dose: Inject 150mg (two 75mg injections) SC every 12 weeks			
	thereafter ☐ Yes or ☐ No: SKYRIZI SELF-INJECTION: Healthcare provider certifies that patient has been trained and is eligible for self-injection					
		☐ Induction	on Dose: To achieve pediatric dose:			
	□ 60kg		Inject 0.75mg/kg			0
☐ STELARA®			0kg - 100kg: Inject 45mg SC 100kg: Inject 90mg SC			0
			ot the contents of 1 prefilled syringe SC on day 1			0
	□ 90mg/ (m) Premied Synnae nor > //0 losi		enance: Inject the contents of 1 prefilled syringe SC on day 29 very 12 weeks thereafter		1	
	☐ Yes or ☐ No: STELARA SELF-INJECTION: Healthcare provider certifies that patient has been trained and is eligible for self-injection			lf-injection		
	□ 100mg/ml Prefilled Syringe		on Dose: Inject 100mg/ml SC at weeks 0 an	d 4	2	0
☐ TREMFYA [™]			intenance: Inject 100mg/ml SC every 8 weeks thereafter			
☐ XELJANZ®	☐ 5mg Tablet	Tablet Take one tablet by mouth twice daily in combination with a nonbiologic Di			60	
☐ XELJANZ [®] XR	□ 11mg Tablet	☐ Take one	tablet by mouth once daily in combination w	ith a nonbiologic DMARD	30	
Signature:	R SIGNATURE: I authorize pharmacy to Da Substitution Permitted rance benefits will be determined by the payor based upon the patient	ate:	Signature: Dispense As \	Written Date:		